



Bivalent vaccine for Newcastle Disease and Infectious Bursal Disease

Description

This vaccine comprises an inactivated combination of the Teso strain of Infectious Bursal Disease (IBD) virus and the Mavoko strain of Newcastle Disease virus, formulated and presented as a water-in-oil emulsion.

Indication

This vaccine is recommended for the vaccination of layers to protect breeder stock and hatched chicks against Newcastle disease and infectious bursal disease (Gumboro disease).

Packages

Vials containing 50 doses and 100 doses.

Dosage and Administration

One dose (0.5 ml) is administered to each bird through subcutaneous injection



(under the skin of the neck) or intramuscularly (in the breast or thigh muscle).

Safety

The bivalent ND and IBD vaccine is safe and does not cause adverse reactions in vaccinated birds. However, any adverse reactions at the site of administration should be reported as soon as they occur

Recommended Vaccination Programme

The vaccine can be administered to birds at around 16-20 weeks of age. For best results, the birds must be primed with live Newcastle disease and Gumboro vaccine 4-6 weeks earlier.

Storage

The vaccine should be stored between +2 and +8 degrees Celsius, away from light, and should not be frozen.



Compiled by: Ogali N.I., Mungube E.O., Mutisya W., Nyambati E.M., Wachira A.M., Ouko V.O and Ngaira V.M

Design and Layout: Odipo S.N

For further information, contact:

Institute Director

Veterinary Research Institute, Muguga,

P.O. Box 32-00902 Kikuyu Email: Director.vsri@kalro.org

Tel No. 0202020572